

EXTRA PAIN RELIEF- acetaminophen, aspirin and caffeine tablet, film coated TIME CAP LABS

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

370R Timely 49483 370 Acetaminophen 250 mg,Aspirin 250 mg (NSAID), and Caffeine 65 mg

DRUG FACTS

Active ingredient (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:

headache

a cold

arthritis

muscular aches

toothache

premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop using and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- you experience any of the following signs of stomach bleeding:

feel faint

have bloody or black stools

vomit blood

have stomach pain that does not get better

ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

DIRECTIONS

Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

OTHER INFORMATION

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, pregelatinized starch, povidone, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? call 1-877-290-4008

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE
IMPRINTED FOIL SEAL OVER THE MOUTH OF THE
BOTTLE IS CUT, TORN, BROKEN OR MISSING



Lot No.:
Exp. Date:

Drug Facts

RETAIN CARTON FOR COMPLETE
PRODUCT INFORMATION

Active ingredient (in each caplet) **Purpose**
Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID) Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches
■ toothache ■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ for more than 10 days for pain unless directed by a doctor ■ for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis or kidney disease ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug or are under a doctor's care for any serious condition

Drug Facts (continued)

Stop using and ask a doctor if ■ symptoms do not improve ■ new symptoms occur ■ pain or fever persists or gets worse ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose,

NDC 49483-370-42
Compare to the active ingredients
in Excedrin® Extra Strength*

timely™

Extra Strength
PAIN RELIEVER
Acetaminophen 250 mg, Aspirin 250 mg (NSAID)
and Caffeine 65 mg
Pain Reliever / Pain Reliever Aid

24 CAPLETS
(Caplet Shaped Tablet)

Actual Size

E3144R

Drug Facts (continued)

mineral oil, pregelatinized starch, povidone, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? call 1-877-298-4008

This product is not manufactured or distributed by
GlaxoSmithKline Pte., owner of the registered
trademark Excedrin® Extra Strength. 3708 1220
Distributed by: Time-Cap Ltd., Inc.
7 Michael Avenue, Farmingdale, NY 11735



NDC 49483-370-42

Extra Strength **PAIN RELIEVER**

*Acetaminophen 250 mg,
Aspirin 250 mg (NSAID),
and Caffeine 65 mg*

Pain Reliever /
Pain Reliever Aid

24 CAPLETS**
(**Capsule-Shaped Tablet)

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: headache, a cold, arthritis, muscular aches, toothache ■ premenstrual and menstrual cramps

Warnings See carton for complete warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling

■ shock ■ asthma (wheezing) **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product **Stomach**

bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning

(anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ **do not use more than directed** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 caplets every 6 hours, not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING** ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Questions or comments? call 1-877-290-4088

Manufactured by: Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale, NY 11735
370R 11/20

Lot No.:

Exp. Date:



NDC 49483-370-20
Compare to the active
ingredients in
Excedrin® Extra Strength†

Extra Strength
PAIN RELIEVER
Acetaminophen 250 mg,
Aspirin 250 mg (NSAID),
and Caffeine 65 mg

Pain Reliever / Fever Reducer

200 CAPLETS**

(**Capsule-Shaped Tablet)

Actual Size



**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE
IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS
CUT, TORN, BROKEN OR MISSING**

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID)* Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses

■ temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ toothache
■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you use:

■ more than 8 caplets in 24 hours, which is the maximum daily amount

■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may

Drug Facts (continued under label)

†This product is not manufactured or distributed by
GlaxoSmithKline plc, owner of the registered trademark
Excedrin® Extra Strength.
370R 1020

Distributed by:

Time-Cap Labs, Inc.

7 Michael Avenue

Farmingdale, NY 11735



1

3

49483-37020

Lot No.:

Exp. Date:



Drug Facts (continued)

cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older ■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of

caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability,

sleeplessness, and, occasionally, rapid heartbeat.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug

contains acetaminophen, ask a doctor or pharmacist.

■ for more than 10 days for pain unless directed by a doctor

■ for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis or kidney disease ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking

any other drug or are under a doctor's care for any serious condition

Stop using and ask a doctor if ■ symptoms do not improve ■ new symptoms occur ■ pain or fever persists or gets worse

■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have

stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product.

If pregnant or breast-feeding, ask a health professional before use. It's especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not use more than directed

■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 2 caplets every 6 hours,

not more than 8 caplets in 24 hours

■ children under 12 years: ask a doctor


Other information

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients: carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, pregelatinized

starch, povidone, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? call 1-877-290-4008



NDC 49483-370-01

Extra Strength
PAIN RELIEVER
*Acetaminophen 250 mg,
 Aspirin 250 mg (NSAID),
 and Caffeine 65 mg*

Pain Reliever / Pain Reliever Aid

100 CAPLETS**
 (**Capsule-Shaped Tablet)

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: headache, a cold, arthritis, muscular aches, toothache, ■ premenstrual and menstrual cramps

Warnings: See carton for complete warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin redness ■ skin edding ■ blisters ■ rash

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen while using this product ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription (NSAIDs) (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during the delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ **do not use more than directed**
 ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information

■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**
 ■ store at 25° C (77°F); excursions permitted between 15°-30° C (59°-86°F)
 ■ use by expiration date on package

Questions or comments? Call 1-877-290-4008

Manufactured by: Timex-Cap Labs, Inc.
 7 Michale Avenue
 Farmingdale, NY 11735

Lot No.: 370R 1020

Exp. Date:

EXTRA PAIN RELIEF

acetaminophen, aspirin and caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-370
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE ((CAPSULE SHAPED TABLET))	Size	18mm
Flavor		Imprint Code	TCL370
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-370-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/16/2021	
2	NDC:49483-370-20	200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2013	
3	NDC:49483-370-00	20000 in 1 CARTON; Type 0: Not a Combination Product	02/22/2016	
4	NDC:49483-370-42	1 in 1 CARTON	07/16/2021	
4		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/01/2013	

Labeler - TIME CAP LABS (037052099)

Registrant - TIME CAP LABS (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABS		037052099	manufacture(49483-370)

Revised: 7/2021

TIME CAP LABS